

Mathematical Analysis of Elliptical Path in the Annulus of Two Circles

Mr Harish Chandra Rajpoot

Feb, 2015

M.M.M. University of Technology, Gorakhpur-273010 (UP), India

1. Introduction:

When a smaller circle exactly lies inside the bigger one (either touching or not touching internally) with their centres separated by a certain distance, then the **locus of the centre of a third tangent (parametric) circle, in the annular region between them, touching the smaller circle externally & the bigger one internally is always an ellipse having the centres of former circles as its foci & its centre at the midpoint of the centres of former circles.** This path becomes a circle if both the smaller & bigger circles are concentric i.e. centres of both the circles, smaller inside bigger, are coincident. (See figure 1 below, showing an elliptical path (dotted curve A'BAB'A') in the annular region between two circles, smaller inside bigger, with their centres separated by a certain distance)

2. Derivation of major axis, minor axis, eccentricity & equation of elliptical path in the annular region between two circles, one inside other, with their centres separated by a certain distance:

Consider a smaller circle, with a radius r , centred at the origin (point O) completely inside the bigger one, with a radius R , centred at the point O' such that distance between their centres O & O' is d . Now consider an **elliptical path** (dotted curve A'BAB'A' with the centre C) traced by the centre of a third (parametric) circle, in the annular region between them, which touches the smaller circle externally & the bigger one internally.

For convenience, let's assume

$$\text{major axis, } AA' = 2a \quad \& \quad \text{minor axis, } BB' = 2b$$

Now, we have

$$OE = OF = r, \quad O'D = O'G = R \quad \& \quad OO' = d$$

$$DE = O'D - O'E = O'D - (O'O + OE) = R - (d + r)$$

$$\therefore A'E = \frac{DE}{2} = \frac{R - (d + r)}{2} = \frac{R - d - r}{2}$$

$$FG = OG - OF = OO' + O'G - OF = d + R - r$$

$$\therefore FA = \frac{FG}{2} = \frac{R + d - r}{2}$$

$$\Rightarrow A'A = A'E + OE + OF + FA = \frac{R - d - r}{2} + r + r + \frac{R + d - r}{2} = \frac{R - d - r + 4r + R + d - r}{2} = R + r$$

$$\therefore \text{Major axis, } 2a = AA' = R + r \quad \dots \dots \dots (I)$$




Figure 1: Elliptical path (dotted curve A'BAB'A' with the centre C) is the locus of the centre of a third tangent (parametric) circle, in the annular region between two circles (smaller inside bigger) with their centres separated by a certain distance d , touching the smaller circle (with the centre O) externally & the bigger one (with the centre O') externally.

Mathematical Analysis of Elliptical Path in the Annulus of Two Circles

Now, join the vertex B of the transverse axis BCB' to the origin O so as the line OB intersects the smaller circle at the point I & then join B to the centre O' & extend the line O'B to intersect the bigger circle at the point H. Now, draw a third tangent circle, with a radius a & centre at the vertex B, which touches the smaller circle externally & the bigger one internally. Thus we have

$$BC = \text{semi minor axis} = b, \quad BI = BH = \text{radius of tangent circle} = a \quad \&$$

$$\Rightarrow OC = OG - CG = (OF + FG) - (CA + AG)$$

$$= (OF + FG) - \left(\frac{AA'}{2} + FA \right) \quad \left(CA = CA' = \frac{AA'}{2} \quad \& \quad AG = FA \right)$$

$$OC = (r + R + d - r) - \left(\frac{R+r}{2} + \frac{R+d-r}{2} \right) = \frac{2R + 2d - 2R - d}{2} = \frac{d}{2}$$

$$\therefore CO' = OO' - OC = d - \frac{d}{2} = \frac{d}{2} \quad \Rightarrow \quad \mathbf{CO = CO' = \frac{d}{2}} \quad \dots \dots \dots (II)$$

Thus, the centre C is the midpoint of line segment OO'. Now in the right triangles BCO & BCO' we have

$$CO = CO' = \frac{d}{2}, \quad BO = BO = b \quad \& \quad \angle BCO = \angle BCO' = 90^\circ$$

Hence, the right triangles BCO & BCO' are congruent.

In right $\triangle BCO$

$$OB = \sqrt{(BC)^2 + (OC)^2} = \sqrt{b^2 + \left(\frac{d}{2}\right)^2} = \sqrt{\frac{4b^2 + d^2}{4}} = \frac{\sqrt{4b^2 + d^2}}{2}$$

$$\Rightarrow OB = OI + IB = r + a \quad \therefore \mathbf{r + a = \frac{\sqrt{4b^2 + d^2}}{2}} \quad \dots \dots \dots (III)$$

but, $OB = O'B = O'H - BH = R - a$ (since, right triangles BCO & BCO' are congruent)

Now, equating the value of OB from the eq(III), we get

$$R - a = r + a \quad \Rightarrow \quad 2a = R - r \quad \text{or} \quad a = \frac{R - r}{2}$$

Now, substituting the value of a in the above eq(III), we get

$$r + \frac{R - r}{2} = \frac{\sqrt{4b^2 + d^2}}{2} \quad \text{or} \quad \frac{\sqrt{4b^2 + d^2}}{2} = \frac{R + r}{2}$$

$$\Rightarrow 4b^2 + d^2 = (R + r)^2 \quad \text{or} \quad 2b = \sqrt{(R + r)^2 - d^2}$$

\therefore Minor axis, $2b = BB' = \sqrt{(R + r)^2 - d^2}$ (IV)

$$\Rightarrow \text{Eccentricity, } e = \sqrt{1 - \frac{b^2}{a^2}} = \sqrt{1 - \left(\frac{2b}{2a}\right)^2} = \sqrt{1 - \left(\frac{\sqrt{(R + r)^2 - d^2}}{R + r}\right)^2}$$

Mathematical Analysis of Elliptical Path in the Annulus of Two Circles

$$= \sqrt{1 - \frac{(R+r)^2 - d^2}{(R+r)^2}} = \sqrt{1 - 1 + \frac{d^2}{(R+r)^2}} = \sqrt{\frac{d^2}{(R+r)^2}} = \frac{d}{R+r}$$

∴ Eccentricity, $e = \frac{d}{R+r}$ ($0 \leq e < 1 \quad \forall R \geq d+r$) (V)

Equation of elliptical path: Since, the elliptical path has its centre at the point $C \left(\frac{d}{2}, 0 \right)$, major axis $2a$ & minor axis $2b$, hence the equation of the elliptical path, assuming centre of the smaller circle as the origin, can be obtained by using **standard formula of ellipse centred at the origin** as follows

$$\frac{X^2}{A^2} + \frac{Y^2}{B^2} = 1 \Rightarrow \frac{\left(x - \frac{d}{2}\right)^2}{a^2} + \frac{(y-0)^2}{b^2} = 1 \quad \text{or} \quad \frac{\left(x - \frac{d}{2}\right)^2}{a^2} + \frac{y^2}{b^2} = 1$$

∴ Equation of elliptical path, $\frac{\left(x - \frac{d}{2}\right)^2}{a^2} + \frac{y^2}{b^2} = 1 \quad \forall a = \frac{R+r}{2} \quad \& \quad b = \sqrt{(R+r)^2 - d^2}$

Above equation is very useful for determining the co-ordinates of any point on the elliptical path & the radius of the parametric circle with centre at any arbitrary point on the elliptical path.

Now, the distance of foci from the centre C of elliptical path, can be determined as follows

$$X = \pm ae \Rightarrow x - \frac{d}{2} = \pm \left(\frac{R+r}{2}\right) \left(\frac{d}{R+r}\right) \quad \text{or} \quad x = \frac{d}{2} \pm \frac{d}{2} \Rightarrow x = 0 \quad \& \quad x = d$$

but $x = 0$ & $x = d$ represents the coordinates of the centres O & O' respectively

Hence, it's obvious from the above values that the centres of two circles are always the foci of the elliptical path in the annulus of these circles, smaller inside bigger, with their centres separated by a certain distance.

3. Radius of the tangent circle with centre at the vertex B: The radius (a) of tangent circle, having centre at the vertex B of the transverse axis BCB' of the elliptical path, touching the smaller circle externally & the bigger one internally (See the above figure 1), is given as

$$a = \frac{R-r}{2} \quad \forall R \geq d+r$$

And **angle α_o** of the line BO joining the centre B of the tangent circle to the origin O with the x-axis (line OO')

In right $\triangle BCO$

$$\tan \angle BOC = \frac{BC}{OC} \Rightarrow \tan \alpha_o = \frac{b}{\left(\frac{d}{2}\right)} = \frac{2 \left(\frac{\sqrt{(R+r)^2 - d^2}}{2} \right)}{d} = \frac{\sqrt{(R+r)^2 - d^2}}{d} = \sqrt{\left(\frac{R+r}{d}\right)^2 - 1}$$

$$\text{or } \tan \alpha_o = \sqrt{\left(\frac{1}{e}\right)^2 - 1} = \sqrt{\frac{1-e^2}{e^2}} = \frac{\sqrt{1-e^2}}{e} \quad \text{or} \quad \alpha_o = \tan^{-1} \left(\frac{\sqrt{1-e^2}}{e} \right)$$

∴ $\alpha_o = \tan^{-1} \left(\sqrt{\left(\frac{R+r}{d}\right)^2 - 1} \right) = \tan^{-1} \left(\frac{\sqrt{1-e^2}}{e} \right) \quad \forall e = \frac{d}{R+r} \quad \& \quad R \geq d+r$

Above, equation can be used to calculate angle α_o for the tangent circle with centre at the vertex B.

4. Derivation of the radius of tangent (parametric) circle touching the smaller circle externally & the bigger one internally:

Consider a tangent (parametric) circle with the centre at any arbitrary point say E on the elliptical path such that it touches the smaller circle at the point D externally & the bigger one at the point F internally. Now join the centre E of the tangent circle to the origin O, centre C & the centre O' by the dotted lines & also draw a perpendicular EM on the x-axis (see the figure 2) such that

$$\angle EOM = \alpha, \angle ECM = \gamma \text{ \& \ } \angle EO'M = \beta$$

Now, the equation of line OE passing through the origin O & inclined at an angle α with the positive direction of x-axis is given as follows

$$y = mx = x \tan \alpha \quad \forall \quad 0 \leq \alpha \leq \pi$$

Now, solving the equation of the line OE & the equation of elliptical path to calculate the co-ordinates of the point of intersection E of the ellipse & the line by substituting $x = y \cot \alpha$ in the equation of elliptical path as follows

$$\frac{\left(x - \frac{d}{2}\right)^2}{a^2} + \frac{y^2}{b^2} = 1 \Rightarrow \frac{\left(y \cot \alpha - \frac{d}{2}\right)^2}{a^2} + \frac{y^2}{b^2} = 1$$

$$\Rightarrow \frac{\left(y \cot \alpha - \frac{d}{2}\right)^2}{\left(\frac{R+r}{2}\right)^2} + \frac{y^2}{\left(\frac{\sqrt{(R+r)^2 - d^2}}{2}\right)^2} = 1 \quad (\text{on substituting the values of } a \text{ \& \ } b)$$

$$\text{or } \left(\frac{\sqrt{(R+r)^2 - d^2}}{2}\right)^2 \left(y \cot \alpha - \frac{d}{2}\right)^2 + \left(\frac{R+r}{2}\right)^2 y^2 = \left(\frac{R+r}{2}\right)^2 \left(\frac{\sqrt{(R+r)^2 - d^2}}{2}\right)^2$$

$$\Rightarrow \left(\frac{(R+r)^2 - d^2}{4}\right) \left(y^2 \cot^2 \alpha + \frac{d^2}{4} - y d \cot \alpha\right) + \frac{(R+r)^2}{4} y^2 = \frac{(R+r)^2}{4} \left(\frac{(R+r)^2 - d^2}{4}\right)$$

$$\Rightarrow \left\{ \left(\frac{(R+r)^2 - d^2}{4}\right) \cot^2 \alpha + \frac{(R+r)^2}{4} \right\} y^2 - \left\{ \left(\frac{(R+r)^2 - d^2}{4}\right) d \cot \alpha \right\} y + \left(\frac{(R+r)^2 - d^2}{4}\right) \frac{d^2}{4} - \frac{(R+r)^2}{4} \left(\frac{(R+r)^2 - d^2}{4}\right) = 0$$

$$\left\{ \frac{(R+r)^2 \operatorname{cosec}^2 \alpha}{4} - \frac{d^2}{4} \cot^2 \alpha \right\} y^2 - \left\{ \left(\frac{(R+r)^2 - d^2}{4}\right) d \cot \alpha \right\} y - \left(\frac{(R+r)^2 - d^2}{4}\right) \left(\frac{(R+r)^2}{4} - \frac{d^2}{4}\right) = 0$$

$$\Rightarrow \left\{ \frac{(R+r)^2 \operatorname{cosec}^2 \alpha}{4} - \frac{d^2}{4} \cot^2 \alpha \right\} y^2 - \left\{ \left(\frac{(R+r)^2 - d^2}{4}\right) d \cot \alpha \right\} y - \left(\frac{(R+r)^2 - d^2}{4}\right) \left(\frac{(R+r)^2}{4} - \frac{d^2}{4}\right) = 0$$

Now, solving above quadratic for the values of y as follows




Figure 2: A tangent (parametric) circle, with the centre at any arbitrary point E on the elliptical path, is considered such that the angle of the line OE with the positive direction of x-axis is α in order to calculate its radius

y

$$\begin{aligned}
 &= \frac{\left(\frac{(R+r)^2 - d^2}{4}\right) dcota \pm \sqrt{\left\{\left(\frac{(R+r)^2 - d^2}{4}\right) dcota\right\}^2 + 4 \left\{\frac{(R+r)^2 \operatorname{cosec}^2 \alpha}{4} - \frac{d^2}{4} \cot^2 \alpha\right\} \left(\frac{(R+r)^2 - d^2}{4}\right)}}{2 \left\{\frac{(R+r)^2 \operatorname{cosec}^2 \alpha}{4} - \frac{d^2}{4} \cot^2 \alpha\right\}} \\
 &= \frac{\left(\frac{(R+r)^2 - d^2}{4}\right) dcota \pm \left(\frac{(R+r)^2 - d^2}{4}\right) \sqrt{d^2 \cot^2 \alpha + (R+r)^2 \operatorname{cosec}^2 \alpha - d^2 \cot^2 \alpha}}{\frac{1}{2} \{(R+r)^2 \operatorname{cosec}^2 \alpha - d^2 \cot^2 \alpha\}} \\
 &= 2 \left(\frac{(R+r)^2 - d^2}{4}\right) \left(\frac{dcota \pm (R+r) \operatorname{cosec} \alpha}{(R+r)^2 \operatorname{cosec}^2 \alpha - d^2 \cot^2 \alpha}\right) = \left(\frac{(R+r)^2 - d^2}{2}\right) \left(\frac{d \cos \alpha \pm (R+r)}{(R+r)^2 - d^2 \cos^2 \alpha}\right) \sin \alpha
 \end{aligned}$$

Taking positive sign, we get

$$\begin{aligned}
 y &= \left(\frac{(R+r)^2 - d^2}{2}\right) \left(\frac{d \cos \alpha + (R+r)}{(R+r)^2 - d^2 \cos^2 \alpha}\right) \sin \alpha \\
 &= \left(\frac{(R+r)^2 - d^2}{2}\right) \left(\frac{d \cos \alpha + (R+r)}{((R+r) + d \cos \alpha)((R+r) - d \cos \alpha)}\right) \sin \alpha = \frac{\sin \alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d \cos \alpha}\right) \\
 \Rightarrow y &= \frac{\sin \alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d \cos \alpha}\right) \geq 0 \quad \forall 0 \leq \alpha \leq \pi \quad \text{hence this value of } y \text{ is accepted}
 \end{aligned}$$

$$\therefore x = y \cot \alpha = \frac{\sin \alpha \cot \alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d \cos \alpha}\right) = \frac{\cos \alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d \cos \alpha}\right)$$

Taking negative sign, we get

$$\begin{aligned}
 y &= \left(\frac{(R+r)^2 - d^2}{2}\right) \left(\frac{d \cos \alpha - (R+r)}{(R+r)^2 - d^2 \cos^2 \alpha}\right) \sin \alpha \\
 &= \left(\frac{(R+r)^2 - d^2}{2}\right) \left(\frac{d \cos \alpha - (R+r)}{((R+r) + d \cos \alpha)((R+r) - d \cos \alpha)}\right) \sin \alpha = \frac{-\sin \alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) + d \cos \alpha}\right) \\
 \Rightarrow y &= \frac{-\sin \alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) + d \cos \alpha}\right) \leq 0 \quad \forall 0 \leq \alpha \leq \pi \quad \text{hence this value of } y \text{ is discarded}
 \end{aligned}$$

Hence, the co-ordinates of the point of intersection E (i.e. the centre of tangent circle) is given as

$$E \equiv (x, y) \equiv \left(\frac{\cos \alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d \cos \alpha}\right), \frac{\sin \alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d \cos \alpha}\right)\right) \quad \forall 0 \leq \alpha \leq \pi$$

Now, in the right $\triangle EMO$, we have

$$OM = x = \frac{\cos \alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d \cos \alpha}\right) \quad \& \quad EM = y = \frac{\sin \alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d \cos \alpha}\right)$$

$$\therefore OE = \sqrt{(OM)^2 + (EM)^2} = \sqrt{\left(\frac{\cos \alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d \cos \alpha}\right)\right)^2 + \left(\frac{\sin \alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d \cos \alpha}\right)\right)^2}$$

Mathematical Analysis of Elliptical Path in the Annulus of Two Circles

$$\Rightarrow OE = \frac{1}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d\cos\alpha} \right) \sqrt{\sin^2\alpha + \cos^2\alpha} = \frac{1}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d\cos\alpha} \right)$$

$$\text{but, } OE = OD + DE = r + ED \Rightarrow ED = OE - r$$

$$\begin{aligned} \therefore ED &= \frac{1}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d\cos\alpha} \right) - r = \frac{1}{2} \left(\frac{(R+r)^2 - d^2 - 2r(R+r) + 2rd\cos\alpha}{(R+r) - d\cos\alpha} \right) \\ &= \frac{1}{2} \left(\frac{R^2 + r^2 + 2rR - d^2 - 2rR - 2r^2 + 2rd\cos\alpha}{(R+r) - d\cos\alpha} \right) = \frac{1}{2} \left(\frac{R^2 - d^2 - r^2 + 2rd\cos\alpha}{(R+r) - d\cos\alpha} \right) \end{aligned}$$

Hence, the radius (R_α) of the tangent (parametric) circle for the given value of the angle α is given as follows

$$R_\alpha = \frac{1}{2} \left(\frac{R^2 - d^2 - r^2 + 2rd\cos\alpha}{(R+r) - d\cos\alpha} \right) \quad \forall 0 \leq \alpha \leq \pi \ \& \ R \geq d + r$$

The above expression is very useful to determine the radius (R_α) of the tangent (parametric) circle for the given value of parametric angle α .

5. Parametric angles β & γ : The parametric angles β & γ are determined for the given value of parametric angle α as follows

In the right $\triangle EMC$, we have

$$\begin{aligned} \tan \angle ECM &= \frac{EM}{CM} = \frac{EM}{OM - OC} \Rightarrow \tan \gamma = \frac{\frac{\sin\alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d\cos\alpha} \right)}{\frac{\cos\alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d\cos\alpha} \right) - \frac{d}{2}} \\ &= \frac{((R+r)^2 - d^2)\sin\alpha}{(R+r)^2\cos\alpha - d^2\cos\alpha - (R+r)d + d^2\cos\alpha} = \frac{((R+r)^2 - d^2)\sin\alpha}{(R+r)^2\cos\alpha - (R+r)d} \end{aligned}$$

$$\therefore \tan \gamma = \frac{((R+r)^2 - d^2)\sin\alpha}{(R+r)^2\cos\alpha - (R+r)d} \quad 0 \leq \gamma \leq \pi \quad \forall 0 \leq \alpha \leq \pi$$

In the right $\triangle EMO'$, we have

$$\begin{aligned} \tan \angle EO'M &= \frac{EM}{O'M} = \frac{EM}{OM - OO'} \Rightarrow \tan \beta = \frac{\frac{\sin\alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d\cos\alpha} \right)}{\frac{\cos\alpha}{2} \left(\frac{(R+r)^2 - d^2}{(R+r) - d\cos\alpha} \right) - d} \\ &= \frac{((R+r)^2 - d^2)\sin\alpha}{(R+r)^2\cos\alpha - d^2\cos\alpha - 2(R+r)d + 2d^2\cos\alpha} = \frac{((R+r)^2 - d^2)\sin\alpha}{(R+r)^2\cos\alpha + d^2\cos\alpha - 2(R+r)d} \end{aligned}$$

$$\therefore \tan \beta = \frac{((R+r)^2 - d^2)\sin\alpha}{((R+r)^2 + d^2)\cos\alpha - 2(R+r)d} \quad 0 \leq \beta \leq \pi \quad \forall 0 \leq \alpha \leq \pi$$

Thus, the parametric angles β & γ can be determined for the given value of angle α . Also these angles can be co-related with each other by using above expressions.

Mathematical Analysis of Elliptical Path in the Annulus of Two Circles

Conclusion: For given two circles, smaller inside the bigger one, with their radii r & R respectively and their centres separated by a certain distance d ($\forall d \geq 0$), the **locus of the centre of a third tangent (parametric) circle** (in the annulus of two circles) touching the smaller circle externally & the bigger one internally, is always an **ellipse** (except circle in case of concentric circles) then all the parameters of this elliptical path are determined as follows

Major axis ($2a$):

$$2a = R + r$$

Minor axis ($2b$):

$$2b = \sqrt{(R + r)^2 - d^2}$$

Eccentricity (e):

$$e = \frac{d}{R + r} \quad (0 \leq e < 1 \quad \forall R \geq d + r)$$

Radius (R_α) of the tangent (parametric) circle: For the given value of the parametric angle α

$$R_\alpha = \frac{1}{2} \left(\frac{R^2 - d^2 - r^2 + 2rd\cos\alpha}{(R + r) - d\cos\alpha} \right) \quad \forall 0 \leq \alpha \leq \pi \quad \& \quad R \geq d + r$$

Parametric angles β & γ :

$$\tan\beta = \frac{((R + r)^2 - d^2)\sin\alpha}{((R + r)^2 + d^2)\cos\alpha - 2(R + r)d} \quad 0 \leq \beta \leq \pi \quad \forall 0 \leq \alpha \leq \pi$$

$$\tan\gamma = \frac{((R + r)^2 - d^2)\sin\alpha}{(R + r)^2\cos\alpha - (R + r)d} \quad 0 \leq \gamma \leq \pi \quad \forall 0 \leq \alpha \leq \pi$$

All the articles above have been derived by the author by using **simple geometry & trigonometry**. All above articles (formula) are very practical & simple to apply in case studies & practical applications of 2-D Geometry. These articles are the most generalised expressions which can be used for the analysis of the elliptical path in the annular region between two circles, smaller inside the bigger one & their centres separated by a certain distance in order to calculate major axis, minor axis, eccentricity & the radius of the tangent (parametric) circle.

Note: Above articles had been derived & illustrated by **Mr H.C. Rajpoot (B Tech, Mechanical Engineering)**

M.M.M. University of Technology, Gorakhpur-273010 (UP) India

Feb, 2015

Email: rajpotharishchandra@gmail.com

Author's Home Page: <https://notionpress.com/author/HarishChandraRajpoot>